

Oral Surgery: Orthodontic Related Procedures (For Ohio Only)

Policy Number: CSDEN328OH.A
 Effective Date: December 1, 2023

[➔ Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Definitions	2
Applicable Codes	3
Description of Services	3
Clinical Evidence	4
U.S. Food and Drug Administration (FDA)	7
References	7
Policy History/Revision Information	8
Instructions for Use	8
Archived Policy Versions	8

Related Dental Policy
<ul style="list-style-type: none"> Medically Necessary Orthodontic Treatment
Related Medical Policy
<ul style="list-style-type: none"> Orthognathic (Jaw) Surgery

Application

This Dental Policy only applies to the state of Ohio. Any requests for services that are stated as unproven or services for which there is a coverage or quantity limit will be evaluated for medical necessity using Ohio Administrative Code 5160-1-01.

Coverage Rationale

Surgical Placement of Temporary Anchorage Device (Not Related to Distraction Osteogenesis or Orthognathic Surgery)

The surgical placement of temporary Anchorage devices are used in conjunction with orthodontic treatment and are indicated for individuals aged 12 and over for the following:

- Intrusion of maxillary teeth
- Molar Distalization
- Canine Retraction and Intrusion Retraction mechanics
- Correction of anterior Open Bite and deep Overbite
- Correction of canted Occlusal Planes
- For other procedures for which anchorage provides the only opportunity for successful completion

The surgical placement of a temporary Anchorage device is not indicated for individuals with any of the following:

- Known allergy to titanium alloy
- Advanced osteoporosis
- Uncontrolled immune or metabolic bone disorders
- Unmanaged medical conditions that result in excessive bleeding, reduced resistance to infection, or poor healing response
- Poor oral hygiene
- Poor quality cortical bone density and volume

Surgical Access of Unerupted Tooth

Surgical access of unerupted tooth is indicated for the following:

- When a normally developing permanent tooth is unable to erupt into a functional position
- For labially impacted teeth if there will be 2-3 mm of gingival cuff present after eruption

Surgical access of unerupted tooth is not indicated for the following:

- For supernumerary teeth
- When surgical access of impacted tooth would threaten vital structures
- Individuals with unmanaged medical conditions that result in excessive bleeding, reduced resistance to infection, or poor healing response

Placement of Device to Facilitate Eruption of Impacted Tooth

This is the placement of an orthodontic bracket, band or other device and attached with a chain, on an unerupted tooth, after surgical exposure, to aid in its eruption. This procedure is done following the surgical access of an unerupted tooth.

Corticotomy (Not Related to Distraction Osteogenesis or Orthognathic Surgery)

Corticotomy (also known as periodontally accelerated osteogenic orthodontics [PAOO] or surgically assisted osteogenic orthodontics [SAOO]) is unproven due to insufficient evidence of efficacy and/or safety.

Mobilization of Erupted or Malpositioned Tooth to Aid Eruption

Mobilization of erupted or malpositioned tooth to aid eruption is indicated for the treatment of ankylosed permanent teeth.

Definitions

Anchorage: Resistance to force. Anchorage may come from any of the following sources: intraoral (teeth, bone, soft tissue, implants), or extraoral (cervical, occipital, cranial) (AAO).

Angle's Classification of Malocclusion: A classification of malocclusion introduced by Edward H. Angle. The governing criterion is the anteroposterior relationship of maxillary and mandibular first molars (AAO).

- Class I Malocclusion (Neuroclusion): A malocclusion in which the mesiobuccal cusp of the maxillary first molar occludes in the buccal groove of the mandibular first molar. "Class I" is sometimes incorrectly used as a synonym for normal occlusion, whereas it signifies only a normal sagittal relationship of maxillary and mandibular posterior teeth as they meet.
- Class II Malocclusion (Distoclusion): A distal (posterior) placement of the mandibular (lower) molar, a mesial (anterior) relationship of the maxillary (upper), or a combination of the two. The mesiobuccal cusp of the maxillary first molar occludes mesial to the buccal groove of the mandibular first molar, usually near the embrasure between the mandibular molar and second premolar.
 - Division 1: A Class II molar relationship with proclined maxillary incisors.
 - Division 2: A Class II molar relationship, usually with the maxillary central incisors tipped lingually, the maxillary lateral incisors tipped labially. This malocclusion, in many instances, has and an excessive Overbite.
 - Subdivision: Subdivision of any malocclusion category denotes a unilateral malocclusion classification (e.g., Class II, division 2, subdivision).
- Class III Malocclusion (Mesioclusion): Mesial (anterior) relationship of the mandibular first molar to the maxillary first molar, a retruded relationship of the maxillary first molar to the mandibular, or a combination of the two. The mesiobuccal cusp of the maxillary first molar will typically occlude near the embrasure between the mandibular first and second molars.

Ankylosis: Abnormal immobility, union or fusion. It may occur between two bones at their articulation (e.g., TMJ) or between teeth and alveolar bone (AAO).

Corticotomy: A surgical procedure that intentionally inflicts mechanical damage on the cortical bone to increase bone remodelling and accelerate the repair and shorten orthodontic treatment time (Fernández-Ferrer).

Distalization: A common descriptor for the biomechanics involved in moving maxillary first and second molars distally and into a Class I molar relationship (AAO).

Intrusion: A translational form of tooth movement directed apically and parallel to the long axis of a tooth (AAO).

Occlusal Plane: The imaginary surface on which upper and lower teeth meet in occlusion. It is actually a compound curved surface, but is commonly approximated by a plane (straight line in the lateral view) based on specific reference points within the dental arches (AAO).

Open Bite: Lack of tooth contact in an occluding position (also called apertognathia) (AAO).

Orthognathic Surgery: Orthognathic Surgery is the surgical correction of abnormalities of the mandible, maxilla, or both. The underlying abnormality may be present at birth or may become evident as the patient grows and develops or may be the result of traumatic injuries. The severity of these deformities precludes adequate treatment through dental treatment alone (AAOMS).

Overbite: Vertical overlap of maxillary teeth over mandibular anterior teeth, usually measured perpendicular to the occlusal plane (AAO).

Retraction: Pertaining to desired posteriorly directed, orthodontic or orthopedic displacements of teeth or of bones of the face (AAO).

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CDT Code	Description
D7280	Exposure of an unerupted tooth
D7282	Mobilization of erupted or malpositioned tooth to aid eruption
D7283	Placement of device to facilitate eruption of impacted tooth
D7292	Placement of temporary anchorage device (screw retained plate)requiring flap
D7293	Placement of temporary anchorage device requiring flap
D7294	Placement of temporary anchorage device without flap
D7296	Corticotomy – one to three teeth or tooth spaces, per quadrant
D7297	Corticotomy – four or more teeth or tooth spaces, per quadrant
D7298	Removal of temporary anchorage device [screw retained plate], requiring flap
D7299	Removal of temporary anchorage device, requiring flap
D7300	Removal of temporary anchorage device without flap
D7997	Appliance removal (not by dentist who placed appliance), includes removal of archbar

CDT® is a registered trademark of the American Dental Association

CPT Code	Description
41899	Unlisted procedure, dentoalveolar structures

CPT® is a registered trademark of the American Medical Association

Description of Services

Temporary Anchorage devices (TADs) are used to improve Anchorage during routine orthodontic therapy. They are gaining popularity and can allow better Anchorage than extraoral headgear which relies on significant patient compliance for success (TADs may also be used for distraction osteogenesis of the mandible and this use is medical in nature and typically covered under the member's medical plan). Impacted teeth are those that are not expected to erupt into their normal position within the

dental arch. The most common impactions occur with third molars and maxillary canine teeth but may involve any teeth. If impacted teeth (other than third molars, primary or supernumerary teeth) are exposed early and there is no tooth size or arch length discrepancies, these teeth will often erupt on their own. However, these teeth may require surgery to expose the tooth and place a bracket, band or other device on the unerupted tooth, after its exposure, to aid eruption. Corticotomy in this context is a relatively new surgical procedure that involves creating cuts in, or removal of alveolar bone for the purpose of accelerating orthodontic treatment.

Clinical Evidence

Temporary Anchorage Devices (TADs)

Antoszewska-Smith et al (2017) conducted a systematic review and meta-analysis to compare the effectiveness of orthodontic miniscrew implants-temporary intraoral skeletal anchorage devices (TISADs)-in anchorage reinforcement during en-masse retraction in relation to conventional methods of anchorage. A search of PubMed, Embase, Cochrane Central Register of Controlled Trials and Web of Science were performed. Relevant articles were assessed for quality according to Cochrane guidelines and the data extracted for statistical analysis. A metaanalysis of raw mean differences concerning anchorage loss, tipping of molars, retraction of incisors, tipping of incisors, and treatment duration was carried out. 14 articles including 616 patients were selected as meeting criteria for detailed analysis, and the quality of the studies was assessed as moderate. Meta-analysis showed that use of TISADs facilitates better anchorage reinforcement compared with conventional methods. On average, TISADs enabled 1.86 mm more anchorage preservation than did conventional methods. The authors concluded that TISADs are more effective than conventional methods of anchorage reinforcement. The average difference of almost 2 mm seems not only statistically but also clinically significant. However, the results should be interpreted with caution because of the moderate quality of the included studies. More high-quality studies on this issue are necessary to enable drawing more reliable conclusions.

Heravi et al (2016) conducted a study evaluating the movement of impacted canines away from the roots of neighboring teeth before full-mouth bracket placement, performed by means of TADs to decrease undesired side effects on adjacent teeth. The study sample consisted of 34 palatally impacted canines, 19 in the experimental group and 15 in the control group. In the experimental group, before placement of brackets, the impacted canine was erupted by means of miniscrews. In the control group, after initiation of comprehensive orthodontics, canine disimpaction was performed by means of a cantilever spring soldered to a palatal bar. At the end of treatment, volume of lateral incisors and canine root resorption were measured and compared by means of a CBCT-derived tridimensional model. Visual Analogue Scale (VAS) score, bleeding on probing (BOP) and gingival index (GI) were recorded. Clinical success rate was also calculated. The volume of root resorption of lateral teeth in the control group was significantly greater than in the experimental group ($p < 0.001$). At the end of treatment, VAS score, GI and BOP were not significantly different between the two groups. From this study, the authors concluded that disimpaction of canines and moving them to the arch can be done successfully carried out with minimal side effects by means of skeletal anchorage.

Manni et al (2016) conducted a study with the aim of evaluating the effectiveness of the treatment of skeletal Class II malocclusions with an acrylic splint Herbst appliance anchored to miniscrews with 2 types of ligation. Sixty patients (mean age, 11.6 years; SD, 1.9) with a bilateral Angle Class II Division 1 malocclusion were retrospectively selected and divided into 3 homogeneous and balanced groups on the basis of the Herbst anchorage used: without anchorage, miniscrews with elastic chains, and miniscrews with metallic ligatures. A cephalometric sagittal occlusion analysis merged with mandibular incisor proclination and skeletal divergence was carried out before and after treatment. To compare the absolute variations within and among the groups, we performed the 1-sample t test for repeated measures and 1-way analysis of variance, respectively. The results showed overjet was reduced similarly in all groups, the mandibular bone base length increased in the group with elastic chains only, and the change in the distance between Point A and pogonion showed the most reduction in the group with elastic chains ($p < 0.05$). Incisive flaring was more pronounced in the group with no anchorage than in the group with elastic chains ($p < 0.001$) and the group with metallic ligatures ($p = 0.003$). The authors concluded that anchorage to miniscrews with elastic chains increases the orthopedic effect of the acrylic splint Herbst appliance and confirmed that skeletal anchorage reduces incisor flaring.

Turkkahraman et al (2016) conducted a controlled study to evaluate the effects of temporary anchorage devices (TADs) in the treatment of skeletal open bites and to compare the results with untreated controls. A total of forty patients with skeletal anterior open bites were assigned to two groups of twenty each. Titanium miniplates fixed bilaterally to the infrazygomatic crest area

were used as TADs and intrusive forces were applied to the posterior teeth with Ni-Ti coil springs. The treatment and normal growth changes were evaluated using 24 measurements (2 angular, 22 linear) with statistically significant differences found between the groups in several of them. In the treatment group, statistically significant upper molar intrusion, posterior rotation of the occlusal plane, anterior rotation of the mandible, and resultant overbite improvement were found. The authors concluded that mild to moderate skeletal anterior open bites could easily be treated with TADs without orthognathic surgery. With the rigid anchorage of miniplates, true molar intrusion of up to 4 mm was achieved. With molar intrusion, anterior rotation of the mandible and a significant reduction in anterior face height were determined.

Lee et al (2015) conducted a clinical study to compare the treatment duration and dentoskeletal changes between two different anchorage systems used to treat maxillary dentoalveolar protrusion and to examine the effectiveness of en-masse retraction using two miniscrews placed in the midpalatal suture. Fifty-seven patients were divided into two groups according to the method of maxillary posterior anchorage reinforcement: midpalatal miniscrews (25 patients, mean age 22 years) and conventional anchorage (32 patients, mean age 19 years). The en-masse retraction period, overall treatment duration, pre-treatment effective ANB angle, and change in the effective ANB angle were compared with an independent-samples t-test. Compared to the headgear group, the duration of en-masse retraction was longer by approximately 4 months in the miniscrew group ($p < 0.001$). However, we found no significant difference in the total treatment duration between the groups. Moreover, a greater change in the effective ANB angle was observed in patients treated with miniscrews than in those treated with the conventional method ($p < 0.05$). The authors concluded that treatment using miniscrews placed in the midpalatal area will allow orthodontists more time to control the anterior teeth during en-masse retraction, without increasing the total treatment duration. Furthermore, it achieves better dentoskeletal control than does the conventional anchorage method, thereby improving the quality of the treatment results.

Bechtold et al (2013) conducted a study to determine the effects of linear force vector(s) from interradicular miniscrews on the distalization pattern of the maxillary arch in adult Class II patients. Twenty-five adult patients with mild to moderate Class II dentition and minimal crowding were collected. Either single (group A) or dual (group B) miniscrews were inserted on the posterior interradicular area to deliver a distalizing force to the main archwire. The displacement patterns of maxillary incisors and molars were measured and compared. Significant distalization in the molars and incisors was shown in both groups. Significantly greater distalization and intrusion of the first molar and intrusive displacement of the incisor, together with significant reduction of the mandibular plane, were noted in group B, in contrast to the rotation of the occlusal plane in group A. The authors concluded that interradicular miniscrews predictably induced total arch distalization, leading to the correction of Class II. Additional miniscrews in the premolar area appear to facilitate intrusion and distalization of the entire arch according to the position of the force vectors.

Xun et al (2013) The aim of this retrospective study was to quantitatively evaluate the treatment effects of intrusion of overerupted maxillary molars using miniscrew implant anchorage and to investigate the apical root resorption after molar intrusion. The subjects included 30 patients whose average ages were 35.5 ± 9.0 years. All patients had received intrusion treatments for overerupted maxillary molars with miniscrew anchorage. There were 38 maxillary first molars and 26 maxillary second molars to be intruded. Two miniscrews were inserted in the buccal and palatal alveolar bone mesial to the overerupted molar. Force of 100-150 g was applied by the elastic chains between screw head and attachment on each side. Lateral cephalograms and panoramic radiographs taken before and after intrusion were used to evaluate dental changes and root resorption of molars. Only 6 of the 128 miniscrews failed. The first and second molars were significantly intruded by averages of 3.4 mm and 3.1 mm respectively ($p < 0.001$). The average intrusion time was more than 6 months. The crown of the molars mesially tilted by averages of 3.1 degrees and 3.3 degrees ($p < 0.001$) for first and second molars. The amounts of root resorption were 0.2-0.4 mm on average. The intrusion treatment of overerupted molars with miniscrew anchorages could be used as an efficient and reliable method to recover lost restoration space for prosthesis. Radiographically speaking, root resorption of molars was not clinically significant after application of intrusive forces of 200 to 300 g.

Corticotomy

A corticotomy performed on the maxilla or mandible is a relatively new procedure designed to decrease orthodontic treatment time. While the overall body of evidence is large, there is a lack of well-designed studies with larger numbers of participants that demonstrate patient selection criteria and the clinical utility of this procedure, as well as the long term effects

In a 2021 systematic review, DE Stefani et al. sought to evaluate corticotomy assisted orthodontic treatment (CAOT) as a treatment for expanding narrow arches in adult orthodontic patients. Six studies met the review criteria. The results showed inconsistent results among studies, and that while CAOT can have better results than conventional palatal expansion, it cannot

be considered alternative treatment for severe posterior cross bite. Additionally, greater predictability is achieved if CAOT is done in conjunction with bone grafting. The authors concluded that the evidence of corticotomy as a treatment for palatal expansion in adult orthodontic treatment has not been well described in literature and only few published reports are available. Further testing is necessary to confirm the validity of this technique.

Apilaomva et al. (2020) conducted a systematic review to evaluate corticotomies effects on the acceleration or facilitation of tooth movements in different orthodontic treatments. Seven randomized controlled clinical trials and two controlled clinical trials were included in the review. In the selected studies the effectiveness of conventional orthodontic treatment was compared with orthodontics assisted by corticotomy or piezocisión. The effect of bone grafts was also evaluated. Variables such as tooth movement, treatment time, bone density and root resorption were studied as well. The methodological quality and evidence of the selected studies was low. Most of the studies observed a statistically significant increase in the rate of dental movement, when performing alveolar corticotomies as an aid in orthodontic treatment; either with the conventional technique or with piezocision. The effect of combining corticotomy with bone grafts was assessed. Corticotomy procedures performed even with conventional methods or piezocision involve a rate increase in dental movement and acceleration during the first months, subsequently returning to baseline values. Bone density may increase as a result of concurrent placement of bone grafting materials during corticotomy procedure. High heterogeneity among studies made it difficult to draw clear conclusions. However, within the limitations of this review, the corticotomy procedures were able to statistically and clinically produce significant temporary decrease in orthodontic tooth movement rate. The available literature about orthodontics facilitated by corticotomy techniques provides low quality evidence, which is why more research is needed. A research with less risk of bias would allow greater comparisons and more significant conclusions.

In a 2018 systematic review, Gil et al. aimed to provide scientific support to validate alveolar corticotomies as a reliable approach to accelerated orthodontics. Three randomized clinical trials, 2 prospective randomized clinical trials, 6 case series and 1 randomized controlled split-mouth study were included. No clinical trials were retrieved. The results showed the mean total treatment time in corticotomy-facilitated orthodontic cases was 8.85 months; control groups treatment duration was 16.4 months. Complications such as pain, swelling, and dentin hypersensitivity were reported. Few studies mentioned patient/clinician satisfaction. The faster and less invasive procedures appeared to be well tolerated. However, the methodological quality of the selected studies was low, with only low to moderate scientific evidence. The authors concluded that corticotomy-facilitated orthodontics resulted in decreased treatment time. Few complications and low morbidity were found. More solid evidence-based research is required to support these results.

Ji et al. (2017) conducted a study to summarize published systematic reviews (SRs) that assess the effects of adjunctive interventions on the acceleration of orthodontic tooth movement (OTM). Electronic and manual searches were performed up to August 2016. Systematic reviews investigating the impact of adjunctive techniques on the promotion of OTM were included. A total of 11 SRs were included in this study. The results showed the quality of evidence ranged from very low to low. The short-term (1-3 months) effects of low-level laser therapy and corticotomy were supported by low-quality evidence. The evidence regarding the efficacy of photo biomodulation, pulsed electromagnetic field, interseptal bone reduction, two vibrational devices (Tooth Masseur and Orthoaccel) and electrical current was of very low quality. Relaxin injections and extracorporeal shock waves were reported to have no impact on OTM according to low- and very low-quality evidence, respectively. Based on currently available information, the authors conclude that low-quality evidence indicates that LLLT and corticotomy are effective to promote OTM in the short term. Future high-quality trials are required to determine the optimal protocols, as well as the long-term effects of LLLT and corticotomy, before warranting recommendations for orthodontics clinics.

Patterson et al. (2016) conducted a systematic review and meta-analysis to examine the evidence for the effectiveness and safety of corticotomy-facilitated orthodontics. Electronic databases were searched for articles that examined the rate of corticotomy-facilitated orthodontic tooth movement and its effects on the periodontium, root resorption, and tooth vitality. Unpublished literature was searched electronically through ClinicalTrials.gov and the ISRCTN registry. Relevant orthodontic journals and reference lists also were checked for eligible studies. Randomized clinical trials (RCTs) and controlled clinical trials (CCTs) were considered. Two article reviewers independently assessed the search results, screened the relevant articles, performed data extraction, and evaluated the methodologic quality of the studies. Fourteen eligible articles (6 RCTs and 8 CCTs) were included in this review. The results showed that there was a statistically meaningful increase in the rate of tooth movement compared with controls for all corticotomy techniques assessed. Some studies reported that acceleration in tooth movement was only temporary (lasting a few months). Corticotomy procedures did not seem to produce unwanted adverse effects on the periodontium, root resorption, and tooth vitality. The quality of the body of evidence was regarded as low owing to the presence of multiple methodologic issues, high risks of bias, and heterogeneity in the included articles. The authors concluded that corticotomy procedures can produce statistically and clinically meaningful temporary increases in the rate of

orthodontic tooth movement with minimal side-effects. Additional high-quality randomized clinical trials are needed to allow more definitive conclusions.

Professional Societies

American Association of Oral and Maxillofacial Surgeons (AAOMS)

In a 2017 Parameter of Care, AAOMS indicates that due to the added advantages of lower costs and morbidity, fast recovery, and decreased duration of orthodontic therapy and decompensation, surgically assisted osteogenic orthodontics (SAOO) may be considered as a treatment option when clinically indicated for pediatric patients requiring distraction osteogenesis orthognathic surgery.

In the same parameter of care, the AAOMS states that mandibular retrusion and maxillary protraction utilizing temporary anchorage devices (TAD) such as screw retained plating systems and miniscrews may be considered as a more conservative, cost efficient, and less morbid treatment alternative to more complex orthognathic surgery for developing skeletal class III malocclusion in pre-and early adolescents.

American Association of Orthodontists (AAO)

In a 2017 clinical practice guideline for orthodontic and dentofacial orthopedics, the AAO states that maxillary and mandibular/dentoalveolar hyper and hypoplasia may require adjunctive anchorage procedures including but not limited to osseointegrated implants, mini-screw implants, miniplates and other temporary anchorage devices

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Temporary anchorage devices are FDA approved for use in patients aged 12 years and older. There are an extensive number of manufacturers of these devices. Refer to the following website for more information and search by specific product name: <http://www.fda.gov/MedicalDevices/default.htm>. (Accessed July 18,2022)

References

- American Association of Oral and Maxillofacial Surgeons 2017 Parameters of Care: Clinical Practice Guidelines for Oral and Maxillofacial Surgery, Surgical Correction of Maxillofacial Skeletal Deformities.
- American Association of Oral and Maxillofacial Surgeons Criteria for Orthognathic Surgery.2020. Available at: https://www.aaoms.org/docs/practice_resources/clinical_resources/ortho_criteria.pdf Accessed June 16, 2022.
- American Association of Orthodontists Clinical Practice Guidelines for Orthodontic and Dentofacial Orthopedics. 2017.
- American Association of Orthodontists AAO Glossary, 2012.
- American Dental Association CDT Codebook 2023.
- Antoszevska-Smith J, Sarul M, Łyczek J, et al. Effectiveness of orthodontic miniscrew implants in anchorage reinforcement during en-masse retraction: A systematic review and meta-analysis. Am J Orthod Dentofacial Orthop. 2017 Mar;151(3):440-455.
- Apalimova A, Roselló À, Jané-Salas E, Arranz-Obispo C, Marí-Roig A, López-López J. Corticotomy in orthodontic treatment: systematic review. Heliyon. 2020;6(5):e04013. Published 2020 May 27. doi:10.1016/j.heliyon.2020.e04013.
- Beadnell, S. Current Therapy in Oral and Maxillofacial Surgery, 1st ed. St. Louis: Mosby 2012. Chapter 16, Management of the Impacted Canine; p.135-145.
- Bechtold TE, Kim JW, Choi TH, et al. Distalization pattern of the maxillary arch depending on the number of orthodontic miniscrews. Angle Orthod. 2013 Mar;83(2):266-73.
- DE Stefani A, Bruno G, Irlandese G, et al. Is the corticotomy assisted orthodontic treatment efficient in the expansion of narrow arches in adult patients? Minerva Dent Oral Sci. 2021 Feb;70(1):44-48.
- Digman S, Hayes S, Niel JL. Oral and Maxillofacial Surgery, 2nd ed. St. Louis: Mosby 2009. Chapter 10, Pediatric Dentoalveolar Surgery; p.165-184.

Fernández-Ferrer L, Montiel-Company JM, Candel-Martí E, et al. Corticotomies as a surgical procedure to accelerate tooth movement during orthodontic treatment: A systematic review. *Med Oral Patol Oral Cir Bucal*. 2016 Nov.

Gil APS, Haas OL Jr, Méndez-Manjón I, et al. Alveolar corticotomies for accelerated orthodontics: A systematic review. *J Craniomaxillofac Surg*. 2018 Mar;46(3):438-445.

Heravi F, Shafae H, Forouzanfar A, et al. The effect of canine disimpaction performed with temporary anchorage devices (TADs) before comprehensive orthodontic treatment to avoid root resorption of adjacent teeth. *Dental Press J Orthod*. 2016 Mar-Apr;21(2):65-72.

Lee, J. *Current Therapy in Oral and Maxillofacial Surgery*, 1st ed. St. Louis: Mosby 2012. Chapter 17, Implants for Orthodontic Anchorage; p.147.

Lee J, Miyazawa K, Tabuchi M, Sato T, et al. Effectiveness of en-masse retraction using midpalatal miniscrews and a modified transpalatal arch: Treatment duration and dentoskeletal changes. *Korean J Orthod*. 2014 Mar; 44(2):88-95.

Manni A, Mutinelli S, Pasini M, et al. Herbst appliance anchored to miniscrews with 2 types of ligation: Effectiveness in skeletal Class II treatment. *Am J Orthod Dentofacial Orthop*. 2016 Jun;149(6):871-80.

Patterson BM, Dalci O, Darendeliler MA, et al. Corticotomies and Orthodontic Tooth Movement: A Systematic Review. *J Oral Maxillofac Surg*. 2016 Mar;74(3):453-73.

Sakar M, Ogle O, Dym H. *Oral and Maxillofacial Surgery*, 2nd ed. St. Louis: Mosby 2009. Chapter 11, Basic and Complex Exodontia and Surgical Management Of Impacted Teeth; p.185-211.

Shroff B, Lindauer S. *Current Therapy in Orthodontics*, 1st ed. St. Louis: Mosby c2010. Chapter 22, Temporary Anchorage Devices: Biomechanical Opportunities and Challenges; p. 278-290.

Turkkahraman H, Sarioglu M. Are temporary anchorage devices truly effective in the treatment of skeletal open bites? *Eur J Dent*. 2016 Oct-Dec; 10(4):447-453.

Wolford L, Stevao E, Alexander C et al. *Peterson's Principles of Oral and Maxillofacial Surgery*, 2nd ed. Hamilton Ontario: BC Decker 2004. Chapter 55, Orthodontics for Orthognathic Surgery; p.1131-1132.

Xun CL, Zhao H, Zeng XL, et al. Intrusion of overerupted maxillary molars with miniscrew implant anchorage: a radiographic evaluation. *J Huazhong Univ Sci Technolog Med Sci*. 2013 Oct; 33(5):780-5.

Yi J, Xiao J, Li H, et al. Effectiveness of adjunctive interventions for accelerating orthodontic tooth movement: a systematic review of systematic reviews. *J Oral Rehabil*. 2017 Aug; 44(8):636-654.

Policy History/Revision Information

Date	Summary of Changes
12/01/2023	New dental policy

Instructions for Use

This Dental Policy provides assistance in interpreting the UnitedHealthcare Community Plan of Ohio dental benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plans may differ. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, please check the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Dental Policy is provided for informational purposes. It does not constitute the practice of medicine or medical advice.

Archived Policy Versions

Effective Date	Guideline Number	Guideline Title
N/A	N/A	N/A